
Development of Leaf Sampling and Interpretation Methods for Almond

Project No.: 07- HORT8-Brown

Project Leader: Patrick Brown
Department of Plant Sciences
University of California - Davis
One Shields Avenue, MS #2
Davis, CA 95616
(530) 752-0929
phbrown@ucdavis.edu

Project Co-Investigator: Jeremy Nunes
Department of Plant Sciences
University of California - Davis
One Shields Avenue, MS#2
Davis, CA 95616
(530) 864-5123
jpnunes@ucdavis.edu

Project Cooperators: John Edstrom, Farm Advisor - Colusa County
Roger Duncan, Farm Advisor - Stanislaus County
Brent Holtz, Farm Advisor - Madera County
Bob Beede, Farm Advisor - Kings County
Almond Board of California

Interpretive Summary:

Ninety percent of growers and consultants participating in the recent CDFA-FREP funded focus groups on Nutrition and subsequent surveys of growers, felt that UC Critical Values (CV's) were not appropriate for current yield levels, were not useful early in the season and did not provide sufficient guidance for nutrient management. Two explanations for this observation are possible, 1) the current CV's are limited in application and are possibly incorrect, or 2) that there are systematic errors in the manner in which critical values are used. While it is not known if UC CV's are incorrect (this will be verified), it is known that they have not been validated for early season use and it is clear that there has been a systematic error in the way leaf sampling and CV's have been used. Currently, standardized leaf samples from random trees scattered through the orchard are collected, analyzed for nutrients and compared with established CV's. If the resultant mean field nutrient concentration is equal or greater than the CV then the field is deemed to be sufficient. In high value crops, however, this is an invalid approach since it will result in half of the field being below the critical value. Growers,

who have observed that a higher 'CV' is beneficial, are in effect, bringing a greater percentage of individual's trees above the CV.

We conclude that the 'problem' with current CV's is not that they are necessarily wrong, but that they do not account for within-field, within-canopy, between season or within-season variability. Preventing the occurrence of a deficiency in any part of an orchard or canopy, at any time of the year, is essential to high productivity and fertilizer use efficiency and is the goal of good growers. Unfortunately, the tools to achieve this economically and in an environmentally sound manner are not available and over-fertilization is currently the only tool growers have to ensure optimal field productivity. The recent CDFA-FREP nutrition focus group demonstrated that growers are aware of this problem and have a clear desire to find a better approach.

This project aims to correct this situation by developing new approaches and interpretation tools that better quantify field and temporal variability, are sensitive to yield and provide for in-season monitoring and fertilizer optimization in almond and pistachio. This project will also offer the unique opportunity to verify current CV's and determine the utility of nutrient ratios as a diagnostic tool.

Objectives:

1. Determine the degree to which leaf nutrient status varies across a range of representative orchards and environments
2. Determine the degree to which nutrient status varies within the canopy and within the year.
3. Validate current CV's and determine if nutrient ratio analysis provides useful information to optimize fertility management.
4. Develop and extend an integrated nutrient BMP for Almond.

Materials and Methods:

All trials have been initiated in 8 or 9 year old microsprinkler irrigated almond orchards of good to excellent productivity planted to Nonpareil (50%) in soils representative of the region and a large percentage of almond acreage. At experiment completion, trees will have reached 11 or 14 yo (after 3 or 5 years), representing their most productive years.

For each of 4 almond sites (Arbuckle, Salida, Madera, Bakersfield), plots are a 10-15 acre contiguous block. Both leaf and nut samples will be collected at 5 times during the season, selected from 114 trees in each plot for a period of 3-5 years. Sample collection will be spaced evenly over time from full leaf expansion to one month post-harvest. As a phenological marker, days past full bloom and stage of nut development will be noted. Light interception, trunk diameter, and individual yields of these trees will also be measured.

Standard leaf sampling protocol will be carried out on exposed, non-fruiting spurs, as well as collecting leaves from fruiting spurs with 1 and multiple fruit to explore different sampling methods. Composite nut samples will be collected from each site. Both leaf and nut samples will be processed by researchers prior to being sent to the DANR Analytical Laboratory located on the UC Davis campus.

Sampling Schedule for Single Site:

| | 1* | 2 | 3 | 4 | 5 | 6** | Total # samples |
|--------------------------------|------------|------------|------------|------------|-----------|-----------|-----------------|
| Sample Type | 6-Apr | 12-May | 16-Jun | 26-Jul | 15-Aug | 15-Sep | |
| Non-fruiting spur leaves | 30 | 114 | 114 | 30 | | 30 | 318 |
| Fruiting spur leaves - 1 nut | 30 | 30 | 30 | 30 | | 30 | 150 |
| Fruiting spur leaves – 2+ nuts | 30 | 30 | 30 | 30 | | 30 | 150 |
| Nut sample | 10 | 10 | 10 | 10 | 10 | *** | 50 |
| Total # samples | 100 | 184 | 184 | 100 | 10 | 90 | 668 |

* Estimated date full leaf expansion

** Estimated harvest

*** Harvest sample

An extensive Grid-Sampling protocol has been established at each site using techniques developed for GIS (with Richard Plant, a leading agronomic statistician). At 54 grid points uniformly distributed across a 10-15 acre block of trees, May and July leaf nutrient status, light interception, trunk diameter and tree yield will be determined in each Nonpareil tree. At 30 of these grid points, the nutrient status and yield of 2 neighboring NP trees will also be collected as independent data points. Initially, non-fruiting spur leaves in exposed positions will be selected for these samples, however, depending on the early results, sampling protocols may be adjusted. Two statistical techniques 'nugget sampling' and 'modified Mantel' statistics will be used, this approach allows for partitioning of variance in nutrient status due to environment, due to genetic variability and 'random' variability and allows for determination of the interactions and dependencies between nutrition and yield and the nature of spatial variability within an orchard.

Overall this experiment will collect far more samples, analyzed for more nutrients (N, K, P, S, Ca, Mg, B, Zn, Mn, Fe), than ever performed before and will collect the individual tree yields associated with each of these samples. This detailed approach is designed to provide the foundation statistical information needed to guide fertilizer practice for the foreseeable future.

Results and Discussion:

All sites have been established and sampling is on schedul and results of the tissue analyses from the DANR laboratory are pending. Once sufficient data are available statistical analysis will begin. Harvest of individual trees is now being planned for as well as the two remaining rounds of leaf and nut sampling.

Recent Publications:

Preparation of publications for this project will commence after data from year one have been compiled and analyzed.